

Ryozo WATANABE\* and Zennoske IWATSUKI\*\*: **New name for  
*Haplocladium* (C. Muell.) C. Muell. (Musci)**

渡辺良象・岩月善之助： 蘚類コバノキヌゴケ属の新学名

As indicated by Index Nominum Genericorum (Farr et al. 1979), the generic name *Haplocladium* (C. Muell.) C. Muell. (1896) is a later homonym of the genus *Haplocladium* Naegeli (1862) in the Rhodophyta. For the moss genus *Haplocladium* (C. Muell.) C. Muell., we propose the following name.

**Bryohaplocladium** R. Watanabe et Z. Iwats., nom. nov.

*Hypnum* subsect. *Haplocladium* C. Muell., Linnaea 42: 459 (1879).

*Haplocladium* (C. Muell.) C. Muell., Nuov. Giorn. Bot. Ital. n. ser. 3: 116 (1896), hom. illeg. [non *Haplocladium* Naegeli, Sitzungsber. Bayer. Akad. Wiss. München 1861 (2): 377 (1862)].

*Rudia* Schimp. in Jaeg., Rev. Bryol. 2: 111 (1875), nom. nud.

Lectotype: *Haplocladium macropilum* C. Muell., fide Britton in N. L. Britton, Fl. Bermuda: 445 (1916). = *H. angustifolium* (Hampe et C. Muell.) Broth. = *Bryohaplocladium angustifolium* (Hampe et C. Muell.) R. Watanabe et Z. Iwats.

**Bryohaplocladium angustifolium** (Hampe et C. Muell.), comb. nov.

*Hypnum angustifolium* Hampe et C. Muell., Bot. Zeit. 13: 88 (1855). = *Haplocladium angustifolium* (Hampe et C. Muell.) Broth. in Engler & Prantl, Nat. Pflanzenfam. 1 (3): 1008 (1907).

**Bryohaplocladium discolor** (Broth. et Par.), comb. nov.

*Thuidium discolor* Broth. et Par., Rev. Bryol. 31: 58 (1904). = *Haplocladium discolor* (Broth. et Par.) Broth. in Engler & Prantl, Nat. Pflanzenfam. 1 (3): 1007 (1907).

**Bryohaplocladium larminatii** (Broth. et Par.), comb. nov.

*Pseudoleskea larminatii* Broth. et Par., Rev. Bryol. 31: 57 (1904). = *Haplocladium larminatii* (Broth. et Par.) Broth. in Engler & Prantl, Nat. Pflanzenfam. ed. 2, 11: 320 (1925).

\* [redacted] Hoya-shi, Tokyo 202. 東京都保谷市 [redacted]

\*\* The Hattori Botanical Laboratory, Obi, Nichinan-shi, Miyazaki-ken 889-25. 服部植物研究所,

**Bryohaplocladium mexicanum** (Card.), comb. nov.

*Haplocladium mexicanum* Card., Rev. Bryol. 36: 110 (1909).

**Bryohaplocladium microphyllum** (Hedw.), comb. nov.

*Hypnum microphyllum* Hedw., Sp. Musc.: 269 (1801). = *Haplocladium microphyllum* (Hedw.) Broth. in Engler & Prantl, Nat. Pflanzenfam. 1 (3): 1007 (1907).

**Bryohaplocladium parvulum** (R. Watanabe), comb. nov.

*Haplocladium parvulum* R. Watanabe in Iwatsuki, Journ. Hattori Bot. Lab. 46: 301 (1979).

**Bryohaplocladium perparaphyllum** (R. Watanabe), comb. nov.

*Haplocladium perparaphyllum* R. Watanabe, Journ. Hattori Bot. Lab. 36: 201 (1972).

**Bryohaplocladium persistens** (C. Muell.), comb. nov.

*Hypnum persistens* C. Muell., Linnaea 43: 485 (1882). = *Haplocladium persistens* (C. Muell.) Broth. in Engler & Prantl, Nat. Pflanzenfam. 1 (3): 1007 (1907).

**Bryohaplocladium porphyropelma** (C. Muell.), comb. nov.

*Haplocladium porphyropelma* C. Muell., Hedwigia 40: 83 (1901).

**Bryohaplocladium schimperi** (Thér.), comb. nov.

*Haplocladium schimperi* Thér., Ann. Crypt. Exot. 3: 75 (1930).

**Bryohaplocladium schwetschkeoides** (Card.), comb. nov.

*Pseudoleskea schwetschkeoides* Card., Bull. Soc. Bot. Genève, ser. 2, 3: 284 (1911). = *Haplocladium schwetschkeoides* (Card.) Broth. in Engler & Prantl, Nat. Pflanzenfam. ed. 2, 11: 320 (1925).

**Bryohaplocladium strictulum** (Card.), comb. nov.

*Thuidium strictulum* Card., Beih. Bot. Centralbl. 17: 29 (1904). = *Haplocladium strictulum* (Card.) Reim., Hedwigia 76: 199 (1937).

**Bryohaplocladium virginianum** (Brid.), comb. nov.

*Hypnum virginianum* Brid., Bryol. Univ. 2: 576 (1827). = *Haplocladium virginianum* (Brid.) Broth. in Engler & Prantl, Nat. Pflanzenfam. 1 (3): 1007 (1907).

The moss genus, *Haplocladium* (C. Muell.) C. Muell., belongs to the Thuidiaceae and is widely distributed in the world. There is no legitimate name to replace *Haplocladium* (C. Muell.) C. Muell., hom. illeg. However, Jaeger (1875) listed *Rudia* Schimper, nom. nud., which included only one species,

*Hypnum fluminense* Hampe, 1875, nom. nud. It, according to Reimers (1937), is a synonym of *Haplocladium microphyllum* (Hedw.) Broth. For this reason, the name *Rudia* Schimp. was considered as a synonym of *Haplocladium* in Index Muscorum, vol. 4 (Wijk et al. 1967). To avoid future confusion, *Bryohaplocladium* appears to be the best name to use in place of *Haplocladium* (C. Muell.) C. Muell.

We thank Prof. M. Chihara of Tsukuba University and Dr. H. Inoue of National Science Museum, Tokyo for informations on *Haplocladium* Naegeli, and also to Dr. A. J. Sharp of University of Tennessee and Dr. S. Hattori of the Hattori Botanical Laboratory for their suggestions.

#### Literature cited

Farr, E. R., J. A. Leussink, and F. A. Stafleu, 1979. Index Nominum Genericorum (Plantarum), vol. 2, 631-1276. Utrecht. Jaeger, A. 1875. Conspectus systematis generum muscorum omnium quem ex operibus et actis bryologorum. Rev. Bryol. 2(8) : 105-112. Reimers, H. 1937. Die europäischen *Haplocladium*-Arten mit besonderer Berücksichtigung ihrer aussereuropäischen Verbreitung und ihrer Verwandtschaft. Hedwigia 76 : 191-298. Wijk, R. van der, W. D. Margadant, and P. A. Florschütz, 1967. Index Muscorum, vol. 4(P-S), 1-604. Utrecht.

\* \* \* \*

コバノキスゴケ属 (*Haplocladium* C. Muell., 1896) はシノブゴケ科 (Thuidiaceae) の藓で, 13種があり, うち一種 *H. angustifolium* はヨーロッパ, アジア, 中南米に広く分布している。この属は東亜でよく分化し, 日本には7種がある。近頃になって, この属には先行名 (*Haplocladium* Naegeli, 1862) が紅藻類にあることが判った。このため, 藓類 *Haplocladium* C. Muell. については新学名, *Bryohaplocladium* R. Watanabe et Z. Iwatsuki を用意した。

□角山栄: 茶の世界史 pp. 225. 1980. 中公新書596. 中央公論社, 東京. ¥440. 茶の飲用について醗酵の紅茶と, 乾燥による緑茶がある。徳川時代のはじめは緑茶がむしろ茶の湯文化と結びついて大いにはやったが, 其後鎮国の間にインド大陸に茶業が進展して紅茶の進出となった。それを日中英の三国を中心として論じたもので中々面白い。近時再び緑茶が文化として見直されつつあるかに見える時に描かれたものであるだけに, 意味が深い。

(前川文夫)